

LBR TP TMDL Modeling Decision Log

(**Note – this document strives to address the major decision points and rationales made during the Model Workgroup and Work Sessions. There are many more discussion points and documents that are available as part of the post meeting summaries, etc. available on the DEQ LBR webpage at:

<http://www.deq.idaho.gov/regional-offices-issues/boise/basin-watershed-advisory-groups/lower-boise-river-wag.aspx>)

- November 28, 2012 thru March 21, 2013 – Monthly Model Workgroup Meetings
 - Discussing mass balance and mechanistic modeling approaches (e.g. Qual2k and AQUATOX)
- March 21, 2013 – Monthly Model Workgroup Meeting
 - AQUATOX is selected to help model TP and periphyton relationships on the LBR
 - Weekly Model Work Sessions established to determine and implement model set-up, calibration, and execution
 - Modeling participants will work cooperatively to “build up” the model by reviewing previously-used model components, including coefficients, parameters, segmentation, time, etc...and determine how to maintain and/or modify the existing model to best meet the needs of the TMDL
 - Although modeling will be a group venture and transparent, DEQ will house and run any model scenarios used in the TMDL due to the regulatory implications
- April 2, 2013 – 1st Weekly Model Work Session
 - A 4-segment unlinked model of the LBR, recently updated by Dr. Richard Park was initially reviewed as a starting point, and to be refined, as appropriate, based on group input, review of current data, appropriateness of segments, and needs of the TMDL, etc.
- April 9, 2013 – 2nd Weekly Model Work Session
 - January 1, 2012 through April 30, 2013 selected as time period for initial model set-up and calibration. These dates were chosen in part to: 1) help capture at least a full year, 2) maximize use of USGS synoptic data (August & October 2012, and March 2013), and 3) allow/verify scour during early in the model simulation (e.g. spring 2012).
 - More discussion centered on the pros/cons of linked vs. unlinked model, and identification of what logical segments for either method should be, appropriate initial conditions, simplifying the model by removing/lumping animals, periphyton, and/or other parameters, etc. – follow-up discussions and decisions to take place at April 16 meeting.
 - Initial modeling timeline discussed and established.
- April 16, 2013 – 3rd Weekly Model Work Session
 - A 9-segment linked model of the LBR will be implemented as the starting point for model set-up and calibration. This was based on input from the group and toward meeting the needs of the TMDL (ability to understand the system as a whole, more easily/functionally run scenarios, etc.)

- However, it was discussed and noted that other potential approaches, including a single-segment approach and sensitivity analyses, may need to be implemented if the 9-segment linked model is not performing well and/or meeting the needs of the TMDL.
- For model simplicity and defensibility, animals will not be included in the initial model set-up and calibration.
- Currently, no changes will be made to the existing periphyton categories; however, the group is taking a closer look to determine if changes should be made.
- Initial conditions selected will represent existing conditions to the extent possible/practicable.
 - It was also discussed that because the model run includes more than a year, the first few months of 2012 (e.g. Jan – April), which will be subsequently modeled again for 2013, may need to be evaluated as an “initial spin-up period,” depending on the model performance.