

## **Temporal and Spatial Nutrient Evolution of the Lower Boise River, Southwest Idaho**

Brian Yelen

MS Hydrologic Sciences Student

Department of Geosciences

Boise State University

[brianyelen@u.boisestate.edu](mailto:brianyelen@u.boisestate.edu)

The Lower Boise River in southwest Idaho becomes increasingly contaminated with nutrients. These increases could be attributed to non-point source contamination from agricultural land use and point source from waste water treatment facilities. Understanding the river chemistry and how it evolves spatially and temporally will help locate the areas that input the highest level of contamination as well as areas that contain the highest concentrations. To determine river chemistry, six separate synoptic sampling events were conducted along the approximately 64 miles from Diversion Dam to the Snake River. These sampling events were conducted seasonally from July 2012 to October 2013. Samples were collected every quarter to half mile with a higher frequency of samples taken between Star Idaho and Caldwell Idaho due to an increased number of inputs from tributaries and drainages. The samples were then analyzed at Boise State University for the anions bromine, chloride, fluoride, nitrate, nitrite, phosphate, and sulfate, using an E.P.A. 300.0 equivalent method. Data from laboratory analysis shows overall trends of increasing concentrations for all anions. Bromine, Fluoride, and Sulfate show similar trends between each sampling event with spikes in concentrations occurring at the same intervals. Nutrient behavior varied leading to questions on the cause of this variation. Point source nutrient spikes were identified at waste water treatment facility outlets to the Lower Boise River, with the greatest spikes at the West Boise facility and the Caldwell facility. The largest unidentified source was a spike located approximately 27 miles east of Diversion Dam. These sampling events were able to complete the annual water chemistry cycle and gain valuable knowledge of the Lower Boise River temporal and spatial nutrient evolution.