



## THE PERFORMANCE OF NATURAL ZEOLITE AS A FEED ADDITIVE IN REDUCING AERIAL AMMONIA AND SLURRY AMMONIUM ION CONCENTRATION IN THE PIG FARM NURSERY

Milić, D., Tofant, A., Vučemilo, M.\*  
Venglovský, J.\*\*, Ondrašovičová, O.\*\*

Pig breeding farm "Dubravica" d. d., Pavla Štoosa 109, 10295 Dubravica  
Croatia

\*Faculty of Veterinary Medicine, University of Zagreb, Heinzelova 55, 10000 Zagreb Croatia

\*\*University of Veterinary Medicine, Komenského 73, 041 81 Košice  
The Slovak Republic

jan@venglovsky.com

### ABSTRACT

The investigations were carried out in a pig nursery on 985 piglets (554 experimental and 431 control piglets) for 50 days. The effect of natural zeolite (55 % clinoptilolite) used as a feed additive on the concentration of ammonia in the air was investigated as an important factor affecting the health and welfare of piglets. Our investigations included the determination of microclimatic parameters, aerial ammonia and physicochemical parameters in the slurry. Results of the study proved that the respective feed additive decreased ammonia emissions in the experimental unit. The addition of 2 % of zeolite to mixed feed resulted in a decrease in ammonia emission by about 33 % and a decrease in ammonium concentration in the slurry by 25 %.

Key words: aerial ammonia; ammonium ion; clinoptilolite; pig slurry; zeolite

### INTRODUCTION

Intensive livestock production is increasingly regarded as an important source of gaseous and malodorous emissions due to the production of large quantities of animal excreta and their composition. One of the important air pollutants is ammonia. Other gases released from animal housings are

methane, carbon dioxide, hydrogen sulphide, nitrous oxide, and additional more than 130 trace gases (7).

The presence of ammonia in animal housings affects the health of housed animals. Ammonia causes direct irritation of the retinal and respiratory mucosa resulting in conjunctivitis, coughing, and reduced lung capacity (9, 14). The possibility of intoxication increases with the duration of stay in an ammonia-rich atmosphere. The maximum allowed concentration of ammonia is 25 ppm.

Efforts aimed at reducing aerial ammonia in animal houses are based on various feed and manure additives, for example zeolites (10, 13), glycocomponents from the *Yucca shidigera* plant (1, 15), and a preparation containing sodium alginate obtained from brown seaweed *Ascophillum nodosum* (12).

Zeolites are naturally occurring three-dimensional, microporous, hydrated aluminosilicate minerals characterized by high internal surface area and high cation exchange capacities. Zeolites remove ammonia from slurry by trapping and exchanging it in its crystalline structure. There are more than 50 different types of natural zeolites differing in their selectivity towards various cations. The zeolite clinoptilolite has a specific affinity for ammonium ions and ammonia (11). Moreover, zeolites are non-toxic, non-hazardous and valuable soil conditioners when added to the soil with manure (2).

In the present study, the effect of a commercial preparation "Pigozen", added to piglet feed, was assessed in terms of aerial ammonia and physicochemical composition (ammonium ion) of the slurry.

## MATERIALS AND METHODS

The study was conducted on Dubravica Pig-breeding farm in Hrvatsko Zagorje, Croatia. Measurements were taken in nursery units during the winter period. Piglets were housed in two equal nurseries, in boxes with 30 animals in each, on a partially slatted floor, and were supplied feed and water *ad libitum*.

There were 431 piglets in the control unit and 554 in the experimental. The experiment lasted 50 days.

The commercial preparation "Pigozen", a natural zeolite containing feed additive, was added to the mixed feed in the experimental group at an amount of 2 % by weight.

The microclimatic parameters (temperature, relative humidity and air velocity) were determined (n = 7) every week using a Testo 625 thermohygrometer and Testo 415 anemometer. CO<sub>2</sub> and NH<sub>3</sub> concentrations were determined by a Dräger-Acuvo gas detector pump with respective detector tubes.

Untreated samples of pig slurry were used as a substrate in the study. Grab-bottle, dark green-brown, malodorous specimens were sampled from the channel under the slatted floor once a week (n = 7).

Standard physicochemical parameters in the slurry (dry matter — DM, inorganic matter — IM, pH, biochemical oxygen demand — BOD<sub>5</sub>, and ammonium, nitrite and nitrate ions) were determined in accordance with standard methods (3, 4) using titration and photometric procedures and a HACH DREL/4000 chemistry/apparatus module.

## RESULTS AND DISCUSSION

The results obtained by the measurement of microclimate parameters in the control and experimental nursery units are presented as means in Table 1.

Table 1. Microclimate parameters in control and experimental nursery units

| Parameter                         | Control unit | Experimental unit |
|-----------------------------------|--------------|-------------------|
| Temperature (°C)                  | 24.6         | 24.2              |
| Relative humidity (%)             | 70.1         | 66.2              |
| Air velocity (m.s <sup>-1</sup> ) | 0.09         | 0.08              |

The piglets' health and productivity are greatly influenced by microclimate parameters. Their values measured met the criteria set for the respective animal species and category and were comparable between the control and experimental units. The mean temperature of about 24 °C corresponded to the requirements for this animal category (20—28 °C). Humidity is a factor closely related to air temperature. The relative humidity ranged from 50 % to 70 % in both control and experimental units. Air velocity ranging from 0.05 to 0.2 m.s<sup>-1</sup> corresponded to the requirements for this animal category

(5, 8). Although natural zeolites have great affinity for small polar molecules like H<sub>2</sub>O, SO<sub>2</sub> and NH<sub>3</sub>, lower relative humidity in experimental unit could not be caused by 2 % addition of zeolites (6).

The effect of "Pigozen" was reflected in a considerable difference in the mean aerial gaseous contamination, especially ammonia, between the control and experimental nursery units (Table 2). According to the manufacturer's instructions, this feed additive greatly inhibits the emission of ammonia from animal excreta, thus improving the housing microclimate, which in turn results in reduced mortality and better health in piglets.

Table 2. Mean level of air pollutants in control and experimental nursery units

| Parameter   | Control unit | Experimental unit |
|---|--------------|-------------------|
| Carbon dioxide CO <sub>2</sub> (% by vol.)                  | 0.16         | 0.14              |
| Ammonia NH <sub>3</sub> (ppm)                               | 4.04         | 2.68              |
| Reduction of NH <sub>3</sub> in comparison with control (%) |              | 33.67             |

During the study period, the concentration of ammonia in the experimental unit was reduced by some 33 % on average, which could be attributed to the effect of clinoptilolite. The concentration of carbon dioxide as an indicator of ventilation showed a negligible decrease in the experimental unit.

The influence of zeolite on physicochemical parameters in the slurry differed according to individual parameters (Table 3 and 4). The pH of slurry showed a very small difference and DM content and IM was about 1 % higher in the experimental group, which is related to zeolite addition. On the other hand, the mean value of BOD<sub>5</sub> was lower in the experimental group.

Table 3. Physicochemical parameters in the slurry from control and experimental nursery unit

| Parameter                                       | Control unit | Experimental unit |
|---|--------------|-------------------|
| pH  | 7.1          | 7.0               |
| Dry matter (%)                                  | 7.1          | 8.1               |
| Inorganic dry matter (%)                        | 24.1         | 25.8              |
| Biochemical oxygen demand (mg.l <sup>-1</sup> ) | 6764         | 6280              |

The addition of zeolite-clinoptilolite to mixed feed reduced the concentration of nitrogen compounds — ammonium and nitrate ions by about 26 % and nitrite ions by 12 % (Table 4).

In conclusion, a positive effect of "Pigozen" was observed on the quality of air in terms of aerial ammonia.

Table 4. Level of nitrogen compounds determined in the slurry from control and experimental nursery unit

| Parameter  | Control unit | Experimental unit |
|--|--------------|-------------------|
| Ammonium $\text{NH}_4^+\text{-N}$ (mg.l <sup>-1</sup> )              | 1880         | 1378              |
| Reduction in $\text{NH}_4^+\text{-N}$ in comparison with control (%) |              | 26.7              |
| Nitrite $\text{NO}_2^-\text{-N}$ (mg.l <sup>-1</sup> )               | 6.4          | 5.6               |
| Reduction of $\text{NO}_2^-\text{-N}$ in comparison with control (%) |              | 12.2              |
| Nitrate $\text{NO}_3^-\text{-N}$ (mg.l <sup>-1</sup> )               | 2385         | 1771              |
| Reduction of $\text{NO}_3^-\text{-N}$ in comparison with control (%) |              | 25.8              |

which was decreased by 33 % in the experimental unit. The concentration of ammonium ions and other investigated nitrogen substances in the slurry was also decreased by between 12 and 26 %.

## REFERENCES

1. Amon, M., Dobeic, M., Misselbrook, T. H., Paln, B. P., Phillips, V. R., Sneath, R. W., 1995: A farm-scale study on the use of De-Odorase® for reducing odour and ammonia emissions from intensive fattening piggeries. *Bioresource Technol.*, 51, 163—169.
2. Andersson, M., 1994: *Performance of Additives in Reducing Ammonia Emissions from Cow Slurry*. Swedish Univ. Of Agric. Sci. Dep. Of Agric. Biosyst. and Technol. Rep., No. 93. JBL Publ., Sweden.
3. Anonymous, 1975: *Standard Methods for the Examination of Water and Wastewater*. 14th edn. APHA-AWWA-WPCF. Springfield.
4. Anonymous, 1992: *Water Analysis Handbook*. HACH Company. Loveland. Colorado.
5. Bähr, H., Schröder, H., Odlm, H. V., 1982: *Bauhygienisch Mezmethoden in angewandte Tierhygiene*. Gustav Fischer Verlag, Jena.
6. Ciambelli, P., Corbo, P., Porcelli, C., Rimoli, A., 1985: Ammonia removal from wastewater by natural zeolites. *Zeolites*, 5, 184—187.
7. Hartung, J., 1995: Effect of aerial pollutants in livestock housing on animal and human health. *2nd Symposium Disinfection, Disinsection, Disinfestation in Animal Health and Environmental Protection*. Umag, September 1995, 79—85.
8. Hartung, J., Phillips, V. R., 1994: Control of gaseous emissions from livestock buildings and manure stores. *J. Agr. Eng. Res.*, 57, 173—189.
9. Hojovec, J., 1994: Possibilities of reducing ammonia emissions from pig houses. *Referatesammlung: 3. Slowenischer Symposium über Umwelthygiene, Desinfektion, Entwesung und Rattenbekämpfung*, 92—95.
10. Krieger, R., Pfeiffer, A., 1995: Prüfung von Futter und Güllezusatzstoffen zur Minderung von Ammoniakemissionen aus Schweineställen. *Dtsch. Tierärztl. Wschr.*, 102, 229—338.
11. Mc Crory, D. F., Hobbs P. J., 2001: Additives to reduce ammonia and odor emissions from livestock wastes. *J. Environ. Qual.*, 30, 345—355.
12. Tofant, A., Vučemilo, M., Barač, I., Mamić, S., 1999: Use of Biopolym® granulate as a feed additive in intensive fattening piggeries. *Proc. of the 11th "in-between" Symposium of ISAH*. Postojna, Slovenia, April 1999, 37.
13. Venglovský, J., Pačajová, Z., Sasaková, N., Vučemilo, M., Tofant, A., 1999: Adsorption properties of natural zeolite and bentonite in pig slurry from the microbiological point of view. *Vet Med.-Czech*, 44, 339—344.
14. Versteegen, M., Tamminga, S., Greers, R., 1994: The effect of gaseous pollutants on animals. In Ap Dewi, I., Axford, R. F. E., Fayeze, I., Marai, M., Omed, H.: *Pollution in Livestock Production Systems*. Cab International, Wallingford, UK, 71—79.
15. Vučemilo, M., Tofant, A., Hadina, S., Barač, D., 2004: Effect of *Yucca schidigera* extract, a feed additive, to reduce air pollutants in pig fattening units. *Proc. of "in-between" Congress of the ISAH: Animal Production in Europe*. Saint-Malo - France, 2004, Vol.1, 83.